Electric Dipole Moments

Wolfgang Altmannshofer
waltmannshofer@perimeterinstitute.ca

PERIMETER INSTITUTE

Fermilab Academic Lectures
“The Allure of Ultrasensitive Experiments”
April 1, 2014

References

M. Pospelov and A. Ritz

Electric dipole moments as probes of new physics

Annals Phys. **318**, 119 (2005) [hep-ph/0504231]

J. Engel, M. J. Ramsey-Musolf and U. van Kolck

Electric Dipole Moments of Nucleons, Nuclei, and Atoms: The Standard Model and Beyond

Prog. Part. Nucl. Phys. **71**, 21 (2013) [arXiv:1303.2371 [nucl-th]]

J. R. Ellis, J. S. Lee and A. Pilaftsis

Electric Dipole Moments in the MSSM Reloaded

JHEP **0810**, 049 (2008) [arXiv:0808.1819 [hep-ph]]

Outline of the Lectures

1. Electric Dipole Moments as Probes of New Physics
   - Introduction
   - EDMs in the Standard Model
   - EDMs Beyond the Standard Model
     - Model Independent Sensitivity to New Physics
     - The SUSY CP Problem

Outline of the Lectures

1. Electric Dipole Moments as Probes of New Physics
   - Introduction
   - EDMs in the Standard Model
   - EDMs Beyond the Standard Model
     - Model Independent Sensitivity to New Physics
     - The SUSY CP Problem
2. EDMs, Flavor Physics, and the Higgs
   - Interplay of EDMs with Flavor Physics
     - Minimal Flavor Violation
     - Generic Flavor Violation
   - Interplay of EDMs with Higgs Physics
     - Higgs CP Properties at the LHC
     - Higgs CP Properties and EDMs

Part I

Electric Dipole Moments as Probes of New Physics

Electric and Magnetic Dipole Moments

interactions of a particle with spin $\vec{S}$
with an electric and magnetic field

$$\mathcal{H} = -\mu \frac{\vec{S}}{|S|} \cdot \vec{B} - d \frac{\vec{S}}{|S|} \cdot \vec{E}$$

electric dipole moment d
magnetic dipole moment μ



Properties under C, P, T

Properties under Charge Conjugation C, Parity P, and Time Reversal T

T :	$\vec{E} \to +\vec{E}$	$\vec{B} \to -\vec{B}$	$\vec{S} \to -\vec{S}$
P :	$\vec{E} \to -\vec{E}$	$\vec{B} \to +\vec{B}$	$\vec{S} \to +\vec{S}$

$$\mathcal{H} = -\mu \frac{\vec{S}}{|S|} \cdot \vec{B} - d \frac{\vec{S}}{|S|} \cdot \vec{E}$$

Properties under C, P, T

Properties under Charge Conjugation C, Parity P, and Time Reversal T

T :	$\vec{E} \to +\vec{E}$	$\vec{B} \to -\vec{B}$	$\vec{S} \to -\vec{S}$
P :	$\vec{E} \to -\vec{E}$	$\vec{B} \to +\vec{B}$	$\vec{S} \to +\vec{S}$

$$\mathcal{H} = -\mu \frac{\vec{S}}{|S|} \cdot \vec{B} - d \frac{\vec{S}}{|S|} \cdot \vec{E}$$



MDMs are P even and T even
EDMs are P odd and T odd

assuming CPT invariance
(= pretty safe assumption):

MDMs are CP conserving
EDMs are CP violating

Relevance for Fundamental Particle Physics

Are EDM measurements accurate enough to probe scales that are relevant for high energy physics? (electro-weak scale, TeV scale)

Relevance for Fundamental Particle Physics

Are EDM measurements accurate enough to probe scales that are relevant for high energy physics? (electro-weak scale, TeV scale)

Typical energy resolution in modern EDM experiments

$$\Delta\text{Energy} \sim 10^{-6}\text{Hz} \sim 10^{-21} eV$$

Relevance for Fundamental Particle Physics

Are EDM measurements accurate enough to probe scales that are relevant for high energy physics? (electro-weak scale, TeV scale)

Typical energy resolution in modern EDM experiments

$$\Delta\text{Energy} \sim 10^{-6}\text{Hz} \sim 10^{-21} eV$$

translates into sensitivities of EDMs at the order of

$$d \lesssim \frac{\Delta\text{Energy}}{\text{electric field}} \sim 10^{-25} ecm \quad , \quad \text{electric field} \sim 10^{4} V/cm$$

Relevance for Fundamental Particle Physics

Are EDM measurements accurate enough to probe scales that are relevant for high energy physics? (electro-weak scale, TeV scale)

Typical energy resolution in modern EDM experiments

$$\Delta\text{Energy} \sim 10^{-6}\text{Hz} \sim 10^{-21} eV$$

translates into sensitivities of EDMs at the order of

$$d \lesssim \frac{\Delta\text{Energy}}{\text{electric field}} \sim 10^{-25} ecm \quad , \quad \text{electric field} \sim 10^{4} V/cm$$

Theoretically inferred scaling of EDMs (see later)

$$d \sim \frac{1}{16\pi^2} \times \frac{1 MeV}{\Lambda^2} \quad \Rightarrow \quad \Lambda \gtrsim 1 TeV$$

Relevance for Fundamental Particle Physics

Are EDM measurements accurate enough to probe scales that are relevant for high energy physics? (electro-weak scale, TeV scale)

Typical energy resolution in modern EDM experiments

$$\Delta\text{Energy} \sim 10^{-6}\text{Hz} \sim 10^{-21} eV$$

translates into sensitivities of EDMs at the order of

$$d \lesssim \frac{\Delta\text{Energy}}{\text{electric field}} \sim 10^{-25} ecm \quad , \quad \text{electric field} \sim 10^{4} V/cm$$

Theoretically inferred scaling of EDMs (see later)

$$d \sim \frac{1}{16\pi^2} \times \frac{1 MeV}{\Lambda^2} \quad \Rightarrow \quad \Lambda \gtrsim 1 TeV$$

EDM experiments are relevant for fundamental particle physics

Relativistic Generalization of EDMs and MDMs

Interaction of a fermion f with the photon field A_μ, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$

$$-d_f \frac{\vec{S}}{|S|} \cdot \vec{E} \qquad \rightarrow$$

$$-\mu_f \frac{\vec{S}}{|S|} \cdot \vec{B} \qquad \rightarrow \qquad e(\bar{f}\gamma_\mu f)A^\mu$$

the usual minimal coupling of fermions with the photon give rise to a magnetic moment with gyromagnetic factor $g = 2$

$$\mu_f = g_f \frac{e}{2m_f}$$

Relativistic Generalization of EDMs and MDMs

Interaction of a fermion f with the photon field A_μ, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$

$$-d_f \frac{\vec{S}}{|S|} \cdot \vec{E} \quad \rightarrow \quad d_f \frac{i}{2} (\bar{f} \sigma_{\mu\nu} \gamma_5 f) F^{\mu\nu}$$

$$-\mu_f \frac{\vec{S}}{|S|} \cdot \vec{B} \quad \rightarrow \quad e(\bar{f} \gamma_\mu f) A^\mu + a_f \frac{e}{4m_f} (\bar{f} \sigma_{\mu\nu} f) F^{\mu\nu}$$

the usual minimal coupling of fermions with the photon give rise to a magnetic moment with gyromagnetic factor $g = 2$

the dimension 5 operators induce an electric dipole moment d_f and an anomalous magnetic moment a_f

$$\mu_f = g_f \frac{e}{2m_f} \quad , \qquad (g_f - 2) = 2a_f$$

d_f and a_f are described by non-renormalizable interactions of fermions with the photon. They are absent for elementary fermions at the classical level, but can be induced by loop corrections.

Electric Dipole Moments in the Standard Model

Sources of CP Violation in the Standard Model

The Standard Model of Particle Physics



CP is violated in nature

$$BR(K_L \to \pi^+\pi^-) \neq 0$$

(Cronin, Fitch 1964)

can be accommodated
in the Standard Model
with 3 generations
$\to$ CKM matrix

Sources of CP Violation in the Standard Model

The Standard Model of Particle Physics



CP is violated in nature

$$BR(K_L \to \pi^+\pi^-) \neq 0$$

(Cronin, Fitch 1964)

can be accommodated
in the Standard Model
with 3 generations
$\to$ CKM matrix

Standard Model
contains another source
of CP violation:
the QCD theta term

more on that later ...

The CKM Matrix

parametrizes the misalignment of up-type quarks and down-type quarks in flavor space

appears in the weak interactions of quarks with the W boson (charged current)



$$V_{\text{CKM}} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix}$$

unitary 3×3 matrix $\rightarrow$ 3 angles 6 phases

5 phases can be reabsorbed by redefinition of the quark fields

$$u \rightarrow e^{i\phi_u} u \quad , \quad d \rightarrow e^{i\phi_d} d \quad , \quad \ldots$$

CKM matrix contains one physical CP violating phase

Tree Level Measurement of the CKM Phase



The CKM phase is O(1)

CKM Phase from Fits of the Unitarity Triangle

Within the experimental and theoretical uncertainties, the CKM matrix gives a consistent description of all observed flavor and CP violating phenomena

Extraction of the CKM phase from the global fit gives

$$\gamma = 69.7^{+1.3}_{-2.8}$$



Quark EDMs from the CKM Matrix

try 1 loop with weak interactions to access the phase of the CKM matrix

$$d_d \propto$$



Quark EDMs from the CKM Matrix

try 1 loop with weak interactions to access the phase of the CKM matrix



$$d_d \propto \frac{e}{16\pi^2}$$

- loop suppressed

Quark EDMs from the CKM Matrix

try 1 loop with weak interactions to access the phase of the CKM matrix

$$d_d \propto \frac{e}{16\pi^2} G_F$$



- loop suppressed
- first order in the weak interactions

Quark EDMs from the CKM Matrix

try 1 loop with weak interactions to access the phase of the CKM matrix

$$d_d \propto \frac{e}{16\pi^2} G_F m_d$$



- loop suppressed
- first order in the weak interactions
- helicity suppressed

Quark EDMs from the CKM Matrix

try 1 loop with weak interactions to access the phase of the CKM matrix

$$d_d \propto \frac{e}{16\pi^2} G_F m_d \mathrm{Im}(V_{td} \quad)$$



- loop suppressed
- first order in the weak interactions
- helicity suppressed
- pick up a CKM element that contains a CP violating phase

Quark EDMs from the CKM Matrix

try 1 loop with weak interactions to access the phase of the CKM matrix

$$d_d \propto \frac{e}{16\pi^2} G_F m_d \mathrm{Im}(V_{td} V_{td}^*) = 0$$



- loop suppressed
- first order in the weak interactions
- helicity suppressed
- pick up a CKM element that contains a CP violating phase
- 1 loop is not sufficient...

Quark EDMs from the CKM Matrix

try 2 loops with weak interactions to access the phase of the CKM matrix

$$d_d \propto \frac{e}{(16\pi^2)^2} G_F^2 m_c^2 m_d$$

$$\times \mathrm{Im}(V_{td} V_{tb}^* V_{cb} V_{cd}^*) \neq 0$$



Quark EDMs from the CKM Matrix

try 2 loops with weak interactions to access the phase of the CKM matrix

$$d_d \propto \frac{e}{(16\pi^2)^2} G_F^2 m_c^2 m_d$$

$$\times \mathrm{Im}(V_{td} V_{tb}^* V_{cb} V_{cd}^*) \neq 0$$



- 2 loop suppressed

Quark EDMs from the CKM Matrix

try 2 loops with weak interactions to access the phase of the CKM matrix



$$d_d \propto \frac{e}{(16\pi^2)^2} G_F^2 m_c^2 m_d$$

$$\times \mathrm{Im}(V_{td} V_{tb}^* V_{cb} V_{cd}^*) \neq 0$$

- 2 loop suppressed
- second order in the weak interactions

Quark EDMs from the CKM Matrix

try 2 loops with weak interactions to access the phase of the CKM matrix



$$d_d \propto \frac{e}{(16\pi^2)^2} G_F^2 m_c^2 m_d$$

$$\times \mathrm{Im}(V_{td} V_{tb}^* V_{cb} V_{cd}^*) \neq 0$$

- 2 loop suppressed
- second order in the weak interactions
- pick up CKM combination with non-zero CP phase

Quark EDMs from the CKM Matrix

try 2 loops with weak interactions to access the phase of the CKM matrix



$$d_d \propto \frac{e}{(16\pi^2)^2} G_F^2 m_c^2 m_d$$

$$\times \mathrm{Im}(V_{td} V_{tb}^* V_{cb} V_{cd}^*) \neq 0$$

- 2 loop suppressed
- second order in the weak interactions
- pick up CKM combination with non-zero CP phase

seems to work!
however when one adds up all 2-loop diagrams one still gets 0...
(Shabalin, 1981)

Quark EDMs from the CKM Matrix

first non-vanishing contribution to quark EDMs arises at the 3-loop level



$$d_d \propto \frac{e}{(16\pi^2)^2} \frac{g_s^2}{16\pi^2} G_F^2 m_c^2 m_d$$

$$\times \mathrm{Im}(V_{td} V_{tb}^* V_{cb} V_{cd}^*) \neq 0$$

- two electro-weak loops
- one additional gluon loop

$$d_d \simeq 10^{-34} ecm$$

(Khriplovich 1986,
Czarnecki, Krause 1997)

Lepton EDMs from the CKM Matrix

for lepton EDMs one needs at least one additional loop
to switch from leptons to quarks and to access the CKM phase
(Khriplovich, Pospelov 1991)



$$d_e \propto \frac{e}{(16\pi^2)^3} \frac{g_s^2}{16\pi^2} G_F^3 m_c^2 m_s^2 m_e$$

$$\times \mathrm{Im}(V_{td} V_{tb^*} V_{cb} V_{cd}^*)$$

- three electro-weak loops
- one additional gluon loop

$$d_e \simeq 10^{-44} ecm$$

(Pospelov, Ritz 2013)

Experimentally Accessible EDMs

- EDMs of paramagnetic systems:
  atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)

Experimentally Accessible EDMs

- EDMs of paramagnetic systems:
  atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)
- EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)

Experimentally Accessible EDMs

- EDMs of paramagnetic systems:
  atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)
- EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)
- EDMs of neutron, proton, deuteron

Experimentally Accessible EDMs

- EDMs of paramagnetic systems:
  atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)
- EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)
- EDMs of neutron, proton, deuteron
- EDM of the muon

Experimentally Accessible EDMs

- EDMs of paramagnetic systems:
  atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)
- EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)
- EDMs of neutron, proton, deuteron
- EDM of the muon

need to predict the EDMs of composite systems
in terms of EDMs of elementary particles

Experimentally Accessible EDMs



CP Violating Interactions

CP-odd Lagrangian at the GeV scale

CP Violating Interactions

CP-odd Lagrangian at the GeV scale

$$\frac{g_s^2}{32\pi^2}\bar{\theta} G^A_{\mu\nu}\tilde{G}^{\mu\nu,\,A}$$

QCD theta term

terms at dimension 4

CP Violating Interactions

CP-odd Lagrangian at the GeV scale

$$\frac{g_s^2}{32\pi^2}\bar{\theta} G^A_{\mu\nu}\tilde{G}^{\mu\nu,\,A}$$

QCD theta term

$$d_f \frac{i}{2}(\bar{f}\sigma_{\mu\nu}\gamma_5 f)F^{\mu\nu}$$

EDMs of quarks and leptons

$$d_q^c \frac{ig_s}{2}(\bar{q}_\alpha \sigma^{\mu\nu} T^A_{\alpha\beta}\gamma_5 q_\beta)G^A_{\mu\nu}$$

chromo EDMs (CEDMs) of quarks

terms at dimension 4 , dimension 5

CP Violating Interactions

CP-odd Lagrangian at the GeV scale

$\frac{g_s^2}{32\pi^2}\bar{\theta} G^A_{\mu\nu}\tilde{G}^{\mu\nu,\,A}$	QCD theta term
$d_f \frac{i}{2}(\bar{f}\sigma_{\mu\nu}\gamma_5 f)F^{\mu\nu}$	EDMs of quarks and leptons
$d_q^c \frac{ig_s}{2}(\bar{q}_\alpha \sigma^{\mu\nu} T^A_{\alpha\beta}\gamma_5 q_\beta)G^A_{\mu\nu}$	chromo EDMs (CEDMs) of quarks
$\frac{W}{3} f^{ABC} G^A_{\mu\nu}\tilde{G}^{\nu\rho,\,B} G^{\mu,\,C}_{\rho}$	Weinberg three gluon operator
$C_{ij}(\bar{f}_i f_i)(\bar{f}_j i\gamma_5 f_j)$	CP violating 4 fermion operators

terms at dimension 4 , dimension 5 , dimension 6, ...

CP Violating Interactions



Calculation of EDMs

EDMs of paramagnetic systems:
atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)

e.g. $|d_{Tl}| \simeq$

Calculation of EDMs

EDMs of paramagnetic systems:
atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)

contain an unpaired electron
$\rightarrow$ mainly sensitive to the electron EDM that sees an enhanced effective electric field inside the atom/molecule

e.g. $|d_{Tl}| \simeq \quad d_e$

Calculation of EDMs

EDMs of paramagnetic systems:
atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)

contain an unpaired electron
$\rightarrow$ mainly sensitive to the electron EDM that sees an enhanced effective electric field inside the atom/molecule

typical enhancement factor for atoms

$$d_{\text{para}} \sim \frac{10\, Z^3 \alpha_{\text{em}}}{J(J+1/2)(J+1)^2}\, d_e \quad ,$$

J: angular momentum
Z: atomic number

for polar molecules the enhancement can be even larger

e.g. $|d_{Tl}| \simeq 585\, d_e$

Calculation of EDMs

EDMs of paramagnetic systems:
atoms (Tl, Fr, ...) and molecules (YbF, ThO, ...)

contain an unpaired electron
$\rightarrow$ mainly sensitive to the electron EDM that sees an enhanced effective electric field inside the atom/molecule

typical enhancement factor for atoms

$$d_{\text{para}} \sim \frac{10\, Z^3 \alpha_{\text{em}}}{J(J+1/2)(J+1)^2}\, d_e \quad ,$$

J: angular momentum
Z: atomic number

for polar molecules the enhancement can be even larger

additional contributions from CP-odd electron nucleon couplings
(induced by CP-odd 4 fermion contact terms)

e.g. $|d_{Tl}| \simeq 585\, d_e + e\ 43\ \text{GeV} \times (C_S^{(0)} - 0.2 C_S^{(1)}) + ...$

Calculation of EDMs

EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)

e.g. $|d_{Hg}| \simeq$

Calculation of EDMs

EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)

all electron spins are paired up
suppressed sensitivity to the electron EDM

e.g. $|d_{Hg}| \simeq 10^{-2}\, d_e$

Calculation of EDMs

EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)

all electron spins are paired up
suppressed sensitivity to the electron EDM

sensitivity to the EDM of the nucleus,
that is mainly induced by CP-odd pion nucleon couplings,
that in turn depend mainly on the quark chromo-EDMs

e.g. $|d_{Hg}| \simeq 10^{-2}\, d_e + 7 \times 10^{-3}\, e\, (\tilde{d}_u - \tilde{d}_d)$

Calculation of EDMs

EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)

all electron spins are paired up
suppressed sensitivity to the electron EDM

sensitivity to the EDM of the nucleus,
that is mainly induced by CP-odd pion nucleon couplings,
that in turn depend mainly on the quark chromo-EDMs

additional contributions from CP-odd electron nucleon couplings
(induced by CP-odd 4 fermion contact terms)

e.g. $|d_{Hg}| \simeq 10^{-2}\, d_e + 7 \times 10^{-3}\, e\, (\tilde{d}_u - \tilde{d}_d) + O(C_S) + ...$

Calculation of EDMs

EDMs of diamagnetic atoms (Hg, Ra, Rn, ...)

all electron spins are paired up
suppressed sensitivity to the electron EDM

sensitivity to the EDM of the nucleus,
that is mainly induced by CP-odd pion nucleon couplings,
that in turn depend mainly on the quark chromo-EDMs

additional contributions from CP-odd electron nucleon couplings
(induced by CP-odd 4 fermion contact terms)

e.g. $|d_{Hg}| \simeq 10^{-2}\, d_e + 7 \times 10^{-3}\, e\, (\tilde{d}_u - \tilde{d}_d) + O(C_S) + ...$

most of the terms come with uncertainties of $>$ O(1)

Calculation of EDMs

proton and neutron EDMs

e.g. $d_n \simeq$

Calculation of EDMs

proton and neutron EDMs

high sensitivity to the constituent quark EDMs

e.g. $d_n \simeq 1.4\,(d_d - \frac{1}{4} d_u)$

Calculation of EDMs

proton and neutron EDMs

high sensitivity to the constituent quark EDMs

additional contributions from CP-odd pion nucleon couplings
(mainly induced by chromo-EDMs)

e.g. $d_n \simeq 1.4\,(d_d - \frac{1}{4} d_u) + 1.1\,e\,(\tilde{d}_d + \frac{1}{2}\tilde{d}_u)$

Calculation of EDMs

proton and neutron EDMs

high sensitivity to the constituent quark EDMs

additional contributions from CP-odd pion nucleon couplings
(mainly induced by chromo-EDMs)

also the Weinberg 3 gluon operator can contribute

e.g. $d_n \simeq 1.4\,(d_d - \frac{1}{4} d_u) + 1.1\,e\,(\tilde{d}_d + \frac{1}{2}\tilde{d}_u) + e\ 22\ \text{MeV}\ w$

Calculation of EDMs

proton and neutron EDMs

high sensitivity to the constituent quark EDMs

additional contributions from CP-odd pion nucleon couplings
(mainly induced by chromo-EDMs)

also the Weinberg 3 gluon operator can contribute

additional contributions from CP-odd 4 fermion operators

e.g. $d_n \simeq 1.4\,(d_d - \frac{1}{4} d_u) + 1.1\, e\, (\tilde{d}_d + \frac{1}{2}\tilde{d}_u) + e\ 22\ \text{MeV}\ w + O(C_{ij}) + \ldots$

Calculation of EDMs

proton and neutron EDMs

high sensitivity to the constituent quark EDMs

additional contributions from CP-odd pion nucleon couplings
(mainly induced by chromo-EDMs)

also the Weinberg 3 gluon operator can contribute

additional contributions from CP-odd 4 fermion operators

e.g. $d_n \simeq 1.4\,(d_d - \frac{1}{4} d_u) + 1.1\,e\,(\tilde{d}_d + \frac{1}{2}\tilde{d}_u) + e\ 22\ \text{MeV}\ w + O(C_{ij}) + \ldots$

most of the terms have uncertainties of O(1)

Current Experimental Bounds

[Griffith et al. 2009 (Hg); Baker et al. 2006 (neutron);
Regan et al. 2002 (Tl); Hudson et al. 2011 (YbF); ACME 2013 (ThO)]

$$d_{Hg} < 3.1 \times 10^{-29}\ ecm$$

$$d_n < 2.9 \times 10^{-26}\ ecm$$

$$|d_e| \leq \left\{ \begin{array}{ll} 1.6 & (Tl) \\ 1.05 & (YbF) \\ 0.089 & (ThO) \end{array} \right\} \times 10^{-27} e\,\mathrm{cm}$$

bounds on the electron EDM assume the absence of contributions from CP-odd electron nucleon couplings

Disentangling Different Contributions



Jung, 2013

different paramagnetic atoms/molecules have different dependence on the electron EDM d_e and the CP-odd electron nucleon interaction C_S.

considering many systems simultaneously allows to bound d_e and C_S separately

SM Predictions for EDMs and the QCD theta term

QCD theta term: $$\frac{g_s^2}{32\pi^2}\bar{\theta}G\tilde{G}$$

the QCD theta term is a dimension 4 operator
$\rightarrow$ not suppressed by any high scale
$\rightarrow$ generically expected to be O(1)

SM Predictions for EDMs and the QCD theta term

QCD theta term: $\frac{g_s^2}{32\pi^2}\bar{\theta}G\tilde{G}$

the QCD theta term is a dimension 4 operator
$\rightarrow$ not suppressed by any high scale
$\rightarrow$ generically expected to be O(1)

also contributes to the EDMs of hadronic systems

$$d_n \sim e\bar{\theta}\frac{m_u m_d}{m_u + m_d}\frac{1}{m_n^2} \sim \bar{\theta} \times 6 \times 10^{-17} ecm$$

experimental bound on d_n translates into the limit: $\theta \lesssim 10^{-9}$

$\rightarrow$ strong CP problem

Dynamical Relaxation of $\bar{\theta}$

add an axion: a pseudoscalar that couples to $G\tilde{G}$

$$\mathcal{L} = \bar{\theta}\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a + \frac{a}{f_a}\frac{g_s^2}{32\pi^2}G\tilde{G}$$

Dynamical Relaxation of $\bar{\theta}$

add an axion: a pseudoscalar that couples to $G\tilde{G}$

$$\mathcal{L} = \bar{\theta}\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a + \frac{a}{f_a}\frac{g_s^2}{32\pi^2}G\tilde{G}$$

$$= \left(\bar{\theta} + \frac{a}{f_a}\right)\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a$$

Dynamical Relaxation of $\bar{\theta}$

add an axion: a pseudoscalar that couples to $G\tilde{G}$

$$\mathcal{L} = \bar{\theta}\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a + \frac{a}{f_a}\frac{g_s^2}{32\pi^2}G\tilde{G}$$

$$= \left(\bar{\theta} + \frac{a}{f_a}\right)\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a$$

below the QCD scale a potential for the axion is induced

$$V(a) \propto \left(\bar{\theta} + \frac{a}{f_a}\right)^2$$

axion acquires a vev: $\langle a \rangle = -\bar{\theta} f_a$

Dynamical Relaxation of $\bar{\theta}$

add an axion: a pseudoscalar that couples to $G\tilde{G}$

$$\mathcal{L} = \bar{\theta}\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a + \frac{a}{f_a}\frac{g_s^2}{32\pi^2}G\tilde{G}$$

$$= \left(\bar{\theta} + \frac{a}{f_a}\right)\frac{g_s^2}{32\pi^2}G\tilde{G} + \frac{1}{2}\partial_\mu a \partial^\mu a$$

below the QCD scale a potential for the axion is induced

$$V(a) \propto \left(\bar{\theta} + \frac{a}{f_a}\right)^2$$

axion acquires a vev: $\langle a \rangle = -\bar{\theta} f_a$

the $G\tilde{G}$ term vanishes in the vacuum
$\rightarrow$ no contribution anymore to the neutron EDM

The Neutron EDM from the CKM Matrix

with the QCD theta term switched off,
the dominant contributions to the neutron EDM in the SM
are not the EDMs of the constituent quarks

$$d_n \simeq \bar{\theta} \times 2.5 \times 10^{-16} ecm + 1.1e(\tilde{d}_d + 0.5\tilde{d}_u) + 1.4(d_d - 0.25 d_u)$$

$+$contributions from the Weinberg operator

$+$contributions from 4 fermion operators

The Neutron EDM from the CKM Matrix

with the QCD theta term switched off,
the dominant contributions to the neutron EDM in the SM
are not the EDMs of the constituent quarks

$$d_n \simeq \bar{\theta} \times 2.5 \times 10^{-16} ecm + 1.1e(\tilde{d}_d + 0.5\tilde{d}_u) + 1.4(d_d - 0.25 d_u)$$

$+$contributions from the Weinberg operator

$+$contributions from 4 fermion operators

the dominant effect arises from a CP violating pion nucleon coupling that is generated at the one loop level from a 4-fermion operator



can be as large as

$$d_n \sim 10^{-31}\ ecm$$

(Khriplovich, Zhitnitski 1982)
(Mannel, Uraltsev, 2012)

The “Electron EDM” from the CKM Matrix

in the Standard Model, the contribution to EDMs of paramagnetic systems from the electron is absolutely negligible ($d_e^{\mathrm{SM}} \sim 10^{-44} ecm$)

$$\text{e.g.} \quad d_{Tl} \simeq -585 d_e - e\ 43\mathrm{GeV} \times (C_S^{(0)} - 0.2 C_S^{(1)})$$

The “Electron EDM” from the CKM Matrix

in the Standard Model, the contribution to EDMs of paramagnetic systems from the electron is absolutely negligible ($d_e^{\mathrm{SM}} \sim 10^{-44} ecm$)

e.g. $d_{Tl} \simeq -585 d_e - e\ 43\mathrm{GeV} \times (C_S^{(0)} - 0.2 C_S^{(1)})$

largest SM contribution comes from the
CP violating electron nucleon couplings



“equivalent electron EDM benchmark”

$d_e^{\mathrm{equiv}} \sim 10^{-39}\ ecm$

(Pospelov, Ritz 2013)

Standard Model Benchmarks for EDMs are Many Orders of Magnitude Below the Current Experimental Sensitivities

$\rightarrow$ EDMs are “Background-Free” Probes of New Physics

EDMs and New Physics

Fundamental CP-odd Phases



EDM operators and $SU(2)_L$ Invariance

EDMs and CEDMs are helicity flipping

$$d_f \frac{i}{2}(\bar{f}\sigma_{\mu\nu}\gamma_5 f)F^{\mu\nu} = d_f \frac{i}{2}(\bar{f}_L\sigma_{\mu\nu}f_R - \bar{f}_R\sigma_{\mu\nu}f_L)F^{\mu\nu}$$

$\rightarrow$ EDM and CEDM operators are not $SU(2)_L$ invariant

EDM operators and $SU(2)_L$ Invariance

EDMs and CEDMs are helicity flipping

$$d_f \frac{i}{2}(\bar{f}\sigma_{\mu\nu}\gamma_5 f)F^{\mu\nu} = d_f \frac{i}{2}(\bar{f}_L\sigma_{\mu\nu}f_R - \bar{f}_R\sigma_{\mu\nu}f_L)F^{\mu\nu}$$

$\rightarrow$ EDM and CEDM operators are not $SU(2)_L$ invariant

above the electro-weak scale, need to add a Higgs doublet to the operators to restore $SU(2)_L$ invariance.

$$H(\bar{f}_L\sigma_{\mu\nu}f_R)F^{\mu\nu}$$

EDM operators and $SU(2)_L$ Invariance

EDMs and CEDMs are helicity flipping

$$d_f \frac{i}{2}(\bar{f}\sigma_{\mu\nu}\gamma_5 f)F^{\mu\nu} = d_f \frac{i}{2}(\bar{f}_L\sigma_{\mu\nu}f_R - \bar{f}_R\sigma_{\mu\nu}f_L)F^{\mu\nu}$$

$\rightarrow$ EDM and CEDM operators are not $SU(2)_L$ invariant

above the electro-weak scale, need to add a Higgs doublet to the operators to restore $SU(2)_L$ invariance.

$$\frac{1}{\Lambda^2}H(\bar{f}_L\sigma_{\mu\nu}f_R)F^{\mu\nu}$$

EDM operators and $SU(2)_L$ Invariance

EDMs and CEDMs are helicity flipping

$$d_f \frac{i}{2}(\bar{f}\sigma_{\mu\nu}\gamma_5 f)F^{\mu\nu} = d_f \frac{i}{2}(\bar{f}_L\sigma_{\mu\nu}f_R - \bar{f}_R\sigma_{\mu\nu}f_L)F^{\mu\nu}$$

$\rightarrow$ EDM and CEDM operators are not $SU(2)_L$ invariant

above the electro-weak scale, need to add a Higgs doublet to the operators to restore $SU(2)_L$ invariance.

$$\frac{1}{\Lambda^2}H(\bar{f}_L\sigma_{\mu\nu}f_R)F^{\mu\nu} \quad \rightarrow \quad \frac{v}{\Lambda^2}(\bar{f}_L\sigma_{\mu\nu}f_R)F^{\mu\nu} \quad , \quad d_f \sim \frac{v}{\Lambda^2}$$

$\rightarrow$ EDMs and CEDMs are secretly dimension 6 operators and decouple with the new physics scale Λ squared

Model Independent Sensitivity to New Physics

example 1: down quark EDM

$$\frac{\tilde{C}}{\Lambda^2} H y_d (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \to \frac{\tilde{C}}{\Lambda^2} m_d \, (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \quad , \quad d_d \sim \frac{\tilde{C} m_d}{\Lambda^2}$$

- Minimal Flavor Violation implies that the down EDM is proportional to the down Yukawa coupling ($\to$ lecture on Tuesday)
- constraint from neutron EDM, assuming $\tilde{C} \sim 1$ with O(1) phase

$$\Lambda \gtrsim 50 \text{ TeV}$$

Model Independent Sensitivity to New Physics

example 1: down quark EDM

$$\frac{\tilde{C}}{\Lambda^2} H y_d (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \to \frac{\tilde{C}}{\Lambda^2} m_d \, (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \quad , \quad d_d \sim \frac{\tilde{C} m_d}{\Lambda^2}$$

- Minimal Flavor Violation implies that the down EDM is proportional to the down Yukawa coupling ($\to$ lecture on Tuesday)
- constraint from neutron EDM, assuming $\tilde{C} \sim 1/16\pi^2$ with O(1) phase

$$\Lambda \gtrsim 5 \text{ TeV}$$

Model Independent Sensitivity to New Physics

example 1: down quark EDM

$$\frac{\tilde{C}}{\Lambda^2} H y_d (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \to \frac{\tilde{C}}{\Lambda^2} m_d \, (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \quad , \quad d_d \sim \frac{\tilde{C} m_d}{\Lambda^2}$$

- Minimal Flavor Violation implies that the down EDM is proportional to the down Yukawa coupling ($\to$ lecture on Tuesday)
- constraint from neutron EDM, assuming $\tilde{C} \sim 1/16\pi^2$ with O(1) phase

$$\Lambda \gtrsim 5 \text{ TeV}$$

example 2: electron EDM

$$\frac{\tilde{C}}{\Lambda^2} H y_e (\bar{e}_L \sigma_{\mu\nu} e_R) F^{\mu\nu} \to \frac{\tilde{C}}{\Lambda^2} m_e \, (\bar{e}_L \sigma_{\mu\nu} e_R) F^{\mu\nu} \quad , \quad d_e \sim \frac{\tilde{C} m_e}{\Lambda^2}$$

- MFV: proportionality to electron Yukawa
- constraint from ThO EDM, assuming $\tilde{C} \sim 1$ with O(1) phase

$$\Lambda \gtrsim 300 \text{ TeV}$$

Model Independent Sensitivity to New Physics

example 1: down quark EDM

$$\frac{\tilde{C}}{\Lambda^2} H y_d (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \to \frac{\tilde{C}}{\Lambda^2} m_d \, (\bar{d}_L \sigma_{\mu\nu} d_R) F^{\mu\nu} \quad , \quad d_d \sim \frac{\tilde{C} m_d}{\Lambda^2}$$

- Minimal Flavor Violation implies that the down EDM is proportional to the down Yukawa coupling ($\to$ lecture on Tuesday)
- constraint from neutron EDM, assuming $\tilde{C} \sim 1/16\pi^2$ with O(1) phase

$$\Lambda \gtrsim 5 \text{ TeV}$$

example 2: electron EDM

$$\frac{\tilde{C}}{\Lambda^2} H y_e (\bar{e}_L \sigma_{\mu\nu} e_R) F^{\mu\nu} \to \frac{\tilde{C}}{\Lambda^2} m_e \, (\bar{e}_L \sigma_{\mu\nu} e_R) F^{\mu\nu} \quad , \quad d_e \sim \frac{\tilde{C} m_e}{\Lambda^2}$$

- MFV: proportionality to electron Yukawa
- constraint from ThO EDM, assuming $\tilde{C} \sim 1/16\pi^2$ with O(1) phase

$$\Lambda \gtrsim 30 \text{ TeV}$$

The Minimal Supersymmetric Standard Model

Supersymmetry (SUSY) implies:
every fermion has a bosonic partner
and vice versa

requires 2 Higgs doublets to give mass
to up-type and down-type fermions

$$\tan\beta = \langle H_u \rangle / \langle H_d \rangle$$

expect at least some SUSY particles
(Higgsinos, stops, gluinos)
at or below O(TeV) for a
natural electro-weak scale





CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

Higgsino and Higgs masses $\mu\tilde{H}_u\tilde{H}_d + B\mu H_u H_d + m^2_{H_u}|H_u|^2 + m^2_{H_d}|H_d|^2$
$\rightarrow$ 2 phases

CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

Higgsino and Higgs masses
$\rightarrow$ 2 phases

$$\mu \tilde{H}_u \tilde{H}_d + B\mu H_u H_d + m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2$$

squark and slepton masses
$\rightarrow$ 15 phases

$$m_Q^2 \tilde{Q}_L^\dagger \tilde{Q}_L + m_U^2 \tilde{U}_R^\dagger \tilde{U}_R + m_D^2 \tilde{D}_R^\dagger \tilde{D}_R$$
$$+ m_L^2 \tilde{L}_L^\dagger \tilde{L}_L + m_E^2 \tilde{E}_R^\dagger \tilde{E}_R$$

CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

Higgsino and Higgs masses
$\rightarrow$ 2 phases

$$\mu\tilde{H}_u\tilde{H}_d + B\mu H_u H_d + m_{H_u}^2|H_u|^2 + m_{H_d}^2|H_d|^2$$

squark and slepton masses
$\rightarrow$ 15 phases

$$m_Q^2\tilde{Q}_L^\dagger\tilde{Q}_L + m_U^2\tilde{U}_R^\dagger\tilde{U}_R + m_D^2\tilde{D}_R^\dagger\tilde{D}_R$$
$$+ m_L^2\tilde{L}_L^\dagger\tilde{L}_L + m_E^2\tilde{E}_R^\dagger\tilde{E}_R$$

gaugino masses
$\rightarrow$ 3 phases

$$m_1\tilde{B}\tilde{B} + m_2\tilde{W}\tilde{W} + m_3\tilde{g}\tilde{g}$$

CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

Higgsino and Higgs masses
$\rightarrow$ 2 phases

$$\mu \tilde{H}_u \tilde{H}_d + B\mu H_u H_d + m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2$$

squark and slepton masses
$\rightarrow$ 15 phases

$$m_Q^2 \tilde{Q}_L^\dagger \tilde{Q}_L + m_U^2 \tilde{U}_R^\dagger \tilde{U}_R + m_D^2 \tilde{D}_R^\dagger \tilde{D}_R$$
$$+ m_L^2 \tilde{L}_L^\dagger \tilde{L}_L + m_E^2 \tilde{E}_R^\dagger \tilde{E}_R$$

gaugino masses
$\rightarrow$ 3 phases

$$m_1 \tilde{B}\tilde{B} + m_2 \tilde{W}\tilde{W} + m_3 \tilde{g}\tilde{g}$$

trilinear couplings
$\rightarrow$ 27 phases

$$A_u\, H_u \tilde{Q}_L^\dagger \tilde{U}_R + A_d\, H_d \tilde{Q}_L^\dagger \tilde{D}_R + A_\ell\, H_d \tilde{L}_L^\dagger \tilde{E}_R$$

CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

Higgsino and Higgs masses
$\rightarrow$ 2 phases

$$\mu \tilde{H}_u \tilde{H}_d + B\mu H_u H_d + m_{H_u}^2 |H_u|^2 + m_{H_d}^2 |H_d|^2$$

squark and slepton masses
$\rightarrow$ 15 phases

$$m_Q^2 \tilde{Q}_L^\dagger \tilde{Q}_L + m_U^2 \tilde{U}_R^\dagger \tilde{U}_R + m_D^2 \tilde{D}_R^\dagger \tilde{D}_R$$
$$+ m_L^2 \tilde{L}_L^\dagger \tilde{L}_L + m_E^2 \tilde{E}_R^\dagger \tilde{E}_R$$

gaugino masses
$\rightarrow$ 3 phases

$$m_1 \tilde{B}\tilde{B} + m_2 \tilde{W}\tilde{W} + m_3 \tilde{g}\tilde{g}$$

trilinear couplings
$\rightarrow$ 27 phases

$$A_u\, H_u \tilde{Q}_L^\dagger \tilde{U}_R + A_d\, H_d \tilde{Q}_L^\dagger \tilde{D}_R + A_\ell\, H_d \tilde{L}_L^\dagger \tilde{E}_R$$

not all phases are physical! (like in the case of the CKM matrix)

CP Violation in the MSSM

The MSSM can contain many new sources of CP violation

Higgsino and Higgs masses
$\rightarrow$ 2 phases

$$\mu\tilde{H}_u\tilde{H}_d + B\mu H_u H_d + m_{H_u}^2|H_u|^2 + m_{H_d}^2|H_d|^2$$

squark and slepton masses
$\rightarrow$ 15 phases

$$m_Q^2\tilde{Q}_L^\dagger\tilde{Q}_L + m_U^2\tilde{U}_R^\dagger\tilde{U}_R + m_D^2\tilde{D}_R^\dagger\tilde{D}_R$$
$$+ m_L^2\tilde{L}_L^\dagger\tilde{L}_L + m_E^2\tilde{E}_R^\dagger\tilde{E}_R$$

gaugino masses
$\rightarrow$ 3 phases

$$m_1\tilde{B}\tilde{B} + m_2\tilde{W}\tilde{W} + m_3\tilde{g}\tilde{g}$$

trilinear couplings
$\rightarrow$ 27 phases

$$A_u\, H_u\tilde{Q}_L^\dagger\tilde{U}_R + A_d\, H_d\tilde{Q}_L^\dagger\tilde{D}_R + A_\ell\, H_d\tilde{L}_L^\dagger\tilde{E}_R$$

not all phases are physical! (like in the case of the CKM matrix)

2 phases can be rotated away...

1-Loop MSSM Contributions to EDMs

Example 1:
Bino-Higgsino loop contribution
to the electron EDM



$d_e \propto$

1-Loop MSSM Contributions to EDMs

Example 1:
Bino-Higgsino loop contribution
to the electron EDM



$$d_e \propto \frac{\alpha_1}{4\pi}$$

- 1-loop suppression

1-Loop MSSM Contributions to EDMs

Example 1:
Bino-Higgsino loop contribution
to the electron EDM



$$d_e \propto \frac{\alpha_1}{4\pi} m_e$$

- 1-loop suppression
- helicity suppression

1-Loop MSSM Contributions to EDMs

Example 1:
Bino-Higgsino loop contribution
to the electron EDM



$$d_e \propto \frac{\alpha_1}{4\pi} m_e \tan\beta$$

- 1-loop suppression
- helicity suppression
- $\tan\beta$ enhancement

1-Loop MSSM Contributions to EDMs

Example 1:
Bino-Higgsino loop contribution
to the electron EDM



$$d_e \propto \frac{\alpha_1}{4\pi} m_e \tan\beta \frac{\mathrm{Im}(\mu m_{\tilde{B}})}{m_{\tilde{e}}^4}$$

- 1-loop suppression
- helicity suppression
- $\tan\beta$ enhancement
- sensitive to the relative phase of the bino and higgsino mass

1-Loop MSSM Contributions to EDMs

Example 2:
gluino loop contribution
to the down quark EDM



$d_d \propto$

1-Loop MSSM Contributions to EDMs

Example 2:
gluino loop contribution
to the down quark EDM



$$d_d \propto \frac{\alpha_s}{4\pi} m_d \tan\beta \frac{\mathrm{Im}(\mu m_{\tilde{g}})}{m_{\tilde{d}}^4}$$

- 1-loop suppression
- helicity suppression
- $\tan\beta$ enhancement
- sensitive to the relative phase of the gluino and higgsino mass

1-Loop MSSM Contributions to EDMs

Example 2:
gluino loop contribution
to the down quark EDM



$$d_d \propto \frac{\alpha_s}{4\pi} m_d \tan\beta \frac{\mathrm{Im}(\mu m_{\tilde{g}})}{m_{\tilde{d}}^4}$$

- 1-loop suppression
- helicity suppression
- $\tan\beta$ enhancement
- sensitive to the relative phase of the gluino and higgsino mass

sensitivity to squarks and sleptons at the level of several TeV to several 10's of TeV (depending on $\tan\beta$)

The SUSY CP Problem



(Hisano @ Moriond EW 2014)

EDM bounds push SUSY particles far above the TeV scale

assumptions:

no cancellations between various contributions

order 1 CP violating phases

The SUSY CP Problem



(Pospelov, Ritz 2005)

possible cancellations can be constrained by looking at several EDM constraints simultaneously

in the plot:

$m_{\text{SUSY}} = 500 GeV$, $\tan\beta = 3$

consider phase of the Higgsino mass θ_μ; and universal phase of all trilinear couplings θ_A

Attempts to Avoid too Large EDMs

decoupling $\leftrightarrow$ small CP phases

Attempts to Avoid too Large EDMs

decoupling $\leftrightarrow$ small CP phases

decouple (some or all) SUSY particles:
split SUSY, or high scale SUSY $\rightarrow$ electro-weak scale is fine tuned

Attempts to Avoid too Large EDMs

decoupling $\leftrightarrow$ small CP phases

decouple (some or all) SUSY particles:
split SUSY, or high scale SUSY $\rightarrow$ electro-weak scale is fine tuned

decouple 1st and 2nd generation sfermions:
effective SUSY $\rightarrow$ compatible with naturalness

Attempts to Avoid too Large EDMs

decoupling $\leftrightarrow$ small CP phases

decouple (some or all) SUSY particles:
split SUSY, or high scale SUSY $\rightarrow$ electro-weak scale is fine tuned

decouple 1st and 2nd generation sfermions:
effective SUSY $\rightarrow$ compatible with naturalness

switch off CP phases using a
smart SUSY breaking mediation mechanism:
pure gauge mediation $\rightarrow$ Higgs mass pushes squarks up to O(10) TeV

Attempts to Avoid too Large EDMs

decoupling $\leftrightarrow$ small CP phases

decouple (some or all) SUSY particles:
split SUSY, or high scale SUSY $\rightarrow$ electro-weak scale is fine tuned

decouple 1st and 2nd generation sfermions:
effective SUSY $\rightarrow$ compatible with naturalness

switch off CP phases using a
smart SUSY breaking mediation mechanism:
pure gauge mediation $\rightarrow$ Higgs mass pushes squarks up to O(10) TeV

+ many other clever constructions that try to suppress CP phases

2-Loop MSSM Contributions to EDMs

in many decoupling frameworks, 2-loop Barr-Zee contributions can still give sizable contributions to EDMs

2-Loop MSSM Contributions to EDMs

in many decoupling frameworks, 2-loop Barr-Zee contributions can still give sizable contributions to EDMs

e.g. stop loops in effective SUSY



$$d_f^{2\text{loop}} \propto \frac{e^4}{(16\pi^2)^2} \frac{m_f}{m_A^2} \frac{\mu A_t}{m_{\tilde{t}}^2}$$

2-Loop MSSM Contributions to EDMs

in many decoupling frameworks, 2-loop Barr-Zee contributions can still give sizable contributions to EDMs

e.g. stop loops in effective SUSY



$$d_f^{2\text{loop}} \propto \frac{e^4}{(16\pi^2)^2} \frac{m_f}{m_A^2} \frac{\mu A_t}{m_{\tilde{t}}^2}$$

e.g. chargino loops in split SUSY



$$d_f^{2\text{loop}} \propto \frac{e^4}{(16\pi^2)^2} \frac{m_f}{m_{\tilde{W}}\mu}$$

2-Loop MSSM Contributions to EDMs

in many decoupling frameworks, 2-loop Barr-Zee contributions can still give sizable contributions to EDMs

e.g. stop loops in effective SUSY



$$d_f^{2\text{loop}} \propto \frac{e^4}{(16\pi^2)^2} \frac{m_f}{m_A^2} \frac{\mu A_t}{m_{\tilde{t}}^2}$$

e.g. chargino loops in split SUSY



$$d_f^{2\text{loop}} \propto \frac{e^4}{(16\pi^2)^2} \frac{m_f}{m_{\tilde{W}} \mu}$$

current bounds on EDMs start to probe the 2 loop contributions if the involved particles are at the TeV scale

Summary

- Electric Dipole Moments are sensitive to CP violation beyond the Standard Model at the TeV scale and beyond
- Standard Model “background” is negligible for the foreseeable future
- EDMs of atoms, molecules, nucleons, etc are sensitive to different combinations of CP violating interactions

  $\rightarrow$ possibility to disentangle the source by measuring many different EDMs