The *Allure* of Natural Neutrinos

Fermilab Academic Lecture Series
Thursday April 24, 2014

Ed Kearns
Boston University

http://www.aspera-eu.org/images/stories/files/Roadmap.pdf



http://www.aspera-eu.org/images/stories/files/Roadmap.pdf

Formaggio and Zeller
10.1103/RevModPhys.84.1307



This lecture will be from an experimental perspective…

- What have we learned from these natural sources of neutrinos already?
- What are the experimental issues?
- What can we learn from them next?

For local interest: can one study these neutrinos with a massive underground LArTPC ?

Neutrino Composition of this Lecture



✸ Cosmological Relic Neutrinos



$m_\nu = 0$ $m_\nu = 1.9\ \mathrm{eV}$

http://physics.aps.org/articles/v3/57
Courtesy of Shankar Agarwal and Hume Feldman, University of Kansas; submitted to Mon. Not. R. Astron. Soc.

✸ Cosmological Relic Neutrinos

$$< T_\nu > = 1.9 \text{ K}$$

$$< E_\nu > = 170 \ \mu\text{eV}$$

$$\sum m_\nu < 0.3 \text{ eV}$$

> 0.05 eV from ν oscillation

$$n_\nu = 168 \text{ cm}^{-3}$$

x 1-100 from gravitational clustering

(Weinberg, 1962) 10.1103/PhysRev.128.1457



FIG. 2. Shape of the upper end of an allowed Kurie plot to be expected in a β^- decay if neutrinos are degenerate up to energy E_F, or in a β^+ decay if antineutrinos are degenerate.

$$\nu_e + {}^3\text{H} \rightarrow {}^3\text{He} + e^-$$

$$T_e = Q_\beta \pm m_\nu$$

+ for relic neutrino detection

– tritium endpoint resolution

$$\sigma_{\text{NCB}}\left({}^3\text{H}\right) v_\nu = (7.84 \pm 0.03) \times 10^{-45} \text{ cm}^2$$

Princeton Tritium Observatory for Light, Early-Universe, Massive-Neutrino Yield

arXiv:1307.47



Fascinating and challenging experimental techniques:

100 g of 3H, large area (graphene substrate)
magnetic trajectory filter (like KATRIN)
RF signal from cyclotron motion (like Project 8)
high resolution calorimeter and TOF
possible annual modulation

Safti, Lisanti, Spitz, Formaggio arXiv:1404.0680v1

STATUS: prototyping the individual techniques

Solar Neutrinos

$$4p \rightarrow {}^{4}\mathrm{He} + 2e^{+} + \gamma + 2\nu_e$$

$$p + p \rightarrow {}^{2}\mathrm{H} + e^{+} + \nu_e$$

$${}^{7}\mathrm{Be} + e^{-} \rightarrow {}^{7}\mathrm{Li} + \nu_e$$

$${}^{8}\mathrm{B} \rightarrow {}^{8}\mathrm{Be} + e^{+} + \nu_e$$



30 years of Ray Davis' Homestake experiment - 800 solar neutrinos



Standard Model

Quarks

J/ψ

τ

W/Z

SN1987a

top quark

neutrino oscillation

Non-zero θ13

Higgs-like boson

30 years of Ray Davis' Homestake experiment - 800 solar neutrinos



Standard Model

Quarks

J/ψ

τ

W/Z

SN1987a

top quark

neutrino oscillation

Non-zero θ13

Higgs-like boson

Kamiokande II and Super-K



neutrino scattering
off atomic electrons

directional detection



Kamiokande II and Super-K



neutrino scattering
off atomic electrons

directional detection



SNO





Should measure here if no ν mixing. Standard Solar Model prediction.

(2001) ν_e *only:* $\Phi_{CC} = 1.76 \pm 0.11$ $\times 10^{-6}$ cm^{-1}s^{-1}

$\nu_e + \nu_\mu + \nu_\tau$: $\Phi_{NC} = 5.09 \pm 0.62$ $\times 10^{-6}$ cm^{-1}s^{-1}


18
16
14
12
10
8
6
4
2
Δm^2 in 10^{-5}eV^2
0.1
0.2
0.3
0.4
0.5
$\sin^2(\theta_{12})$
KamLAND
Solar ν
Experiments

Solar neutrinos have delivered:

- CC – NC smoking gun
- θ_{12}, Δm_{12} same as KamLAND
- ν_1 ν_2 ordering (mass hierarchy)
- General picture of solar cycle (*pp*, *pep*, ^{7}Be, ^{8}B)

Indications…

- day/night asymmetry from matter effect in the earth

Not yet:

- Spectral distortion of ^{8}B
- *hep* neutrinos
- CNO neutrinos (competing detailed solar models)


Spectral distortion expected
KamLAND
Solar ν
Experiments
Day/night asymmetry expected
18
16
14
12
10
8
6
4
2
Δm^2 in $10^{-5}eV^2$
0.1
0.2
0.3
0.4
0.5
$\sin^2(\theta_{12})$

$$A_{\mathrm{DN}}^{\mathrm{fit}} = [-3.2 \pm 1.1(\mathrm{stat}) \pm 0.5(\mathrm{syst})]\% \qquad 2.7\sigma$$





a tiny tension


SK I Spectrum
Neutrino Spectrum Sys. Uncertainty
Energy Resolution Sys. Uncertainty
Energy Scale Sys. Uncertainty
Data/MC (unoscillated)
E_{kin} in MeV
SK II Spectrum
Data/MC (unoscillated)
E_{kin} in MeV
SK III Spectrum
MSG applied
Data/MC (unoscillated)
E_{kin} in MeV
SK IV Spectrum
Data/MC (unoscillated)
E_{kin} in MeV


P_{ee}^{day}
$P_{ee}^{d}(E_\nu) = c_\circ + c_1(E_\nu - 10) + c_2(E_\nu - 10)^2$
SK only
Solar+KamLAND$_{exp}$
SK+SNO
Solar global$_{exp}$
SNO only
0.0
0.1
0.2
0.3
0.4
0.5
0.6
3
4
5
6
7
8
9
10
11
12
13
14
15
E_ν in MeV


1.0
0.8
0.6
0.4
0.2
0.0
P
E
$1 - \frac{1}{2}\sin^2\theta_{12}$
Quasi-vacuum oscillation
Matter effect conversion
$\sin^2\theta_{12}$


P_{ee} versus ν Energy
arXiv:1307.7335
0.8
0.6
0.4
0.2
0
P_{ee}
SK/SNO ^{8}B
solar+KamLAND
solar
Super-K
Borexino ^{7}Be
Borexino pep
Borexino ^{8}B
Combined pp
Combined CNO
10^{-1}
1
10
ν Energy

Solar outlook

- Hyper-K: bigger, but shallower, and probably lower photocoverage. $E_{thresh} \approx 7$ MeV

- Borexino and SNO+ will try for pp, pep, CNO

- LArTPC?

5 MeV threshold
spallation background needs study
photon trigger required

electron energy ∝ Eν
~ 100 events per day (34 kton)

$$\nu_e + {}^{40}\mathrm{Ar} \rightarrow {}^{40}\mathrm{K}^* + e^-$$



ICARUS (simulation)

✸ Terrestrial Antineutrinos

http://arxiv.org/pdf/1403.4072v1.pdf



Study heat production in the Earth's interior

Terrestrial Antineutrinos have been observed





✸ Supernova Neutrinos







✸ Supernova Neutrinos



- ★ Guaranteed signal – if you run long enough.
- ★ Enormous statistics in a megaton-scale detector.
- ★ Early warning before light and directional pointing
- ★ Time profile and spectra of great astrophysical interest. Possibilities such as Si-burning and black hole formation.
- ★ Standard picture: Initial burst of ν_e and cooling tail of equal flavors
- ★ Matter effects in SN and in earth may be revealed.
- ★ May reveal fundamental neutrino physics as well.

Supernova Rate



Milky Way: few percent chance per year of spectacular signal

500 kton detector adds 30-50 event burst from Andromeda (M31)

Also a search strategy of time coincidence of 1-2 events from more distant galaxies

Supernova Observables



Time profile

Energy Spectrum

Event Rates in Hyper-K



Entire inner volume (0.74 Mton) should be useable for SN burst.

165K – 230K IBD events

7K – 8K ES events

Enough Statistics to Distinguish SN Models





Directional Pointing







Accuracy of
1-2 degrees

Mass Hierarchy Determination

Neutrino – antineutrino matter resonance swaps with Normal – Inverted Hierarchy





Size of neutronization burst may suggest hierarchy

Observation of shock wave in IBD events favors IH

LArTPC?

- Complementary to WC (ν_e not anti-ν_e)
- 1000's of events in 34kt
- Good energy correlation
- Photon trigger?
  Or trigger by nearby modest sized WC?



Channel	Events *Livermore* model	Events *GKVM* model
$\nu_e + ^{40}\mathrm{Ar} \rightarrow e^- + ^{40}\mathrm{K}^*$	2308	2848
$\overline{\nu}_e + ^{40}\mathrm{Ar} \rightarrow e^+ + ^{40}\mathrm{Cl}^*$	194	134
$\nu_x + e^- \rightarrow \nu_x + e^-$	296	178
Total	**2794**	**3160**

Diffuse Relic Supernova Neutrinos



search in this energy range

- ★ Integrated SN neutrino flux from all galaxies (to $z \approx 1$)
- ★ Probes star formation models as well as supernova models.
- ★ **Not yet observed!** Should be observable by megaton-scale WC detector.
- ★ Daunting for even a massive LArTPC. Spallation background rates need study.





Remarkable improvement with Gd tagging of neutron.

Hyper-K LOI arXiv:1109.3262v1

LArTPC? See challenges for solar neutrinos.
Roughly 40 events in a 34 kt x 10 year exposure

✸ Atmospheric Neutrinos






Super–K ν_μ
Fréjus ν_μ
Fréjus ν_e
AMANDA ν_μ
unfolding
forward folding
IceCube ν_μ
unfolding
forward folding
conventional ν_μ
conventional ν_e
This Work ν_e
prompt ν_μ, ν_e
$E_\nu^2 \Phi_\nu$ [GeV cm^{-2}s^{-1}sr^{-1}]
$\log_{10}$ (E_ν [GeV])





- Exponential energy spectrum energies 100 MeV – 10 TeV
- Flux ratios (well estimated and self-calibrated)
  - mu/e
  - up/down
- Pathlength: 15-13000 km
- Varying matter density





Super-K I+II+III+IV Combined Dataset



Until recently, atmospheric neutrinos provided tightest constraint on θ_{23}



Atmospheric neutrinos have delivered:

- Discovery of neutrino oscillation
- θ_{23}, Δm^2_{23} same values as long-baseline experiments
- Oscillation pattern
- ν_τ appearance (3.8σ)

Not yet:

- Independent measurement of θ_{13}
- Mass hierarchy
- Octant of θ_{23}
- CP violation δ


FAR
DETECTOR
Cosmic Ray Proton
muon neutrinos
Accelerator Proton


$P(\nu_\mu \to \nu_\mu)$
T2K
MINOS/NOvA
LBNE
Cosine Zenith Angle
Energy [GeV]


$P(\nu_\mu \to \nu_e)$
T2K
MINOS/NOvA
LBNE
Cosine Zenith Angle
Energy [GeV]

Oscillograms: Graphical representations of neutrino mixing probability



Resonance in ν_μ–ν_e for NH

…

(in antinu for IH)

First Octant minus Second Octant





θ_{23} = **0.4** vs. θ_{23} = 0.5

θ_{23} = **0.6** vs. θ_{23} = 0.5

In 1st Octant:
More multi-GeV muon survival
Less electron appearance

Results: using reactor θ_{13} constraint





- Both free and constrained fits prefer 2nd octant
- 1.2σ preference for inverted hierarchy
  sensitivity is 0.9σ

} Not significant!

Hyper-Kamiokande



Sensitivity studies simply scale SK result to large exposure, i.e. assume the same detector performance





Octant and CP-δ dependency







ICAL @ INO (Iron Calorimeter at India-based Neutrino Observatory)







- 50 kton mass
- 29000 RPCs
- 1.3T magnetic field
- Best acceptance for vertical muons
- 2-3σ MH in 5-10 years

http://dx.doi.org/10.1016/j.nima.2012.03.017

PINGU (Precision IceCube Next Generation Upgrade)





+ Enormous statistics (25K evt/yr)
+ Full containment of 10 GeV νμCC
+ Possible ν/antiν by $d\sigma/dy$ (ArXiv:1303.0758)
+ Recent work includes shower (CC nu e)
− No up/down normalization
− Zenith acceptance & detector response may be challenging to control

PINGU LOI arXiv:1401.2046v1



Use IceCube and DeepCore as an active veto



3-4 Mton effective volume

10 GeV neutrinos – ideal for core resonance

PINGU LOI arXiv:1401.2046v1



(a) Track-like events.



(b) Cascade-like events.

3σ for MH in < 3 years





LArTPC



- High resolution:
- NC BG rejection
- Direction/energy (see all charged)
- ν/anti-ν handles
- Above are needed to compensate for modest mass
- Magnetize?





LBNE Sensitivity Studies via Performance Estimates

	Lepton (electron)	Lepton (muon)	Hadronic System
Angular Resolution	2°	2°	10°
Energy Resolution	$10\% / \sqrt{E}$	2% (FC) 15% (PC)	$30\% / \sqrt{E}$

A. Guglielmi, Neutrino 2010, Ghandi et al., hep-ph/0807.2759.



A. Blake, LBNE

LBNE study	Neutrino (true)	Antineutrino (true)	Tag method(s)
ν_e-like	30%	6%	p ID only
Anti ν_e-like	43%	22%	p ID only
ν_μ-like	27% / 57%	5% / 0%	p ID / μ-e decay
Anti ν_μ-like	49% / 19%	19% / 24%	p ID / μ-e decay


Ratio to No Oscillations
1
0.5
0
Statistical Uncertainty
1
10
10^2
10^3
10^4
Reconstructed L_ν / E_ν (km/GeV)


Data/Prediction (null oscillation)
1.8
1.6
1.4
1.2
1
0.8
0.6
0.4
0.2
0
Super-K Preliminary
240.4 kt-yr
1
10
10^2
10^3
10^4
L/E (km/GeV) (SK 1+2+3+4)



Arguably – 34 kton LArTPC is more interesting than 500 kton WC!


Atmospheric Neutrinos
LAr Detector Simulation
340 kt-yrs
Normal Hierarchy
sin²θ23=0.6
sin²θ23=0.5
sin²θ23=0.4
Mass Hierarchy
Determination
Inverted Hierarchy
sin²θ23=0.6
sin²θ23=0.5
sin²θ23=0.4
Sensitivity (√Δχ²)
10
5
0
0
0.5
1
1.5
0
0.5
1
1.5
2
δCP / π
δCP / π

High Energy Cosmic Neutrinos




Background Atmospheric Muon Flux
Bkg. Atmospheric Neutrinos (π/K)
Background Stat. and Syst. Uncertainties
Atmospheric Neutrinos (90% CL Charm Limit)
Signal+Bkg. Best-Fit Astrophysical E^{-2} Spectrum
Data
PRELIMINARY
60 TeV
Events per 988 Days
10^2
10^1
10^0
10^{-1}
10^2
10^3
Deposited EM-Equivalent Energy in Detector (TeV)



http://arxiv.org/abs/1402.0467v2





Fig. 5. Sky map in equatorial coordinates of the TS value from the maximum likelihood point source analysis. The most significant cluster consists of five events—all showers and including the second highest energy event in the sample—with a final significance of 8%. This is not sufficient to identify any neutrino sources from the clustering study. The galactic plane is shown as a curved gray line with the galactic center at the bottom left denoted by a filled gray square. Best-fit locations of individual events (listed in Table 1) are indicated with vertical crosses (+) for showers and angled crosses (×) for muon tracks.

IceCube Collaboration, DOI: 10.1126/science.1242856

Neutrino Composition of Neutrino 2014 Boston – See you again soon!



http://neutrino2014.bu.edu/

Registration fee goes up May 1! Hotels are pricey and filling up!