
Fossils, Genes,
and
The Origin of Organs



modified from Radinsky 1987

EXTANT

EXTINCT



EXTANT

EXTINCT



EXTANT

EXTINCT

Expeditions targeted to key parts of the tree of life



Developmental genetics of non-model organisms

Expeditions targeted to key parts of the tree of life

EXTANT

EXTINCT



Developmental genetics of non-model organisms

Expeditions targeted to key parts of the tree of life

EXTANT

EXTINCT




Tioga Co.
Lycoming Co.
PENNSYLVANIA
Devonian
N
1840
site
15
Blossburg
N
Powys Curve
1995
site
15


Highlands
Tidal
Flats
Sea
O
SC
V
W
S

CW

Science
2 April 2004
Vol. 304 No. 5667
Pages 1–152 $10
AAAS


A.
b1
entc
h.r
B.
a1
e
entc
ect.r
ectc
C.
a2
d
c
b2
a3
entc
h.r
rad.f
ul.f
ectc
D.
a2
b2
c
rad.f
ul.f
ectc
E.
h.r
f
g
h
ectc
F.
b1
entc



modified from Radinsky 1987


Flat, wide head
Dorsal eyes
modified from Radinsky 1987


Neck
modified from Radinsky 1987



modified from Radinsky 1987

SECOND EDITION

EVOLUTION OF THE EARTH

ROBERT H. DOTT, JR.
ROGER L. BATTEN



R. Dott and R. Batten,
EVOLUTION OF THE EARTH
2nd edition



R. Dott and R. Batten,
EVOLUTION OF THE EARTH
2nd edition

Arctic Islands unexplored



East Greenland well studied

Ongoing Catskill Project

R. Dott and R. Batten,
EVOLUTION OF THE EARTH
2nd edition

BULLETIN OF CANADIAN PETROLEUM GEOLOGY
VOL. 24, NO. 4 (DECEMBER 1976), P. 485-639

THE MIDDLE-UPPER DEVONIAN CLASTIC WEDGE OF THE FRANKLINIAN GEOSYNCLINE

ASHTON EMBRY[1] AND J. EDWARD KLOVAN[2]

[1], 335 - 8th Avenue S.W., Calgary, Alberta.
nt of Geology, University of Calgary, Calgary, Alberta.

Composition — Fine-grained sandstones in the Fram Fm are quartz arenites and quartzose lithic arenites (Plate 24, Fig. 4). The average composition of five fine-grained sandstones from the type section is 70 ± 3 per cent quartz; 1 ± 1 per cent feldspar; 5 ± 2 per cent rock fragments (mainly shale); 6 ± 2 per cent chert; 12 ± 3 per cent clay (mainly kaolinite, minor illite and chlorite); 3 ± 3 per cent calcite; 2 ± 2 per cent porosity. Biotite and muscovite are also present and are usually concentrated in dark laminations along with carbonaceous material. Heavy minerals are rare and consist mainly of tourmaline, zircon, rutile and garnet. Most grains are subround.

Part of the quartz content is authigenic cement. The clay is interpreted to be mainly of diagenetic origin and tends to occur in the coarser sandstones. Sparry calcite is clearly a cement and is most common in very fine grained sandstones and coarse siltstones. The samples point counted had a fairly low calcite content (max. 9 per cent) but others range as high as 30 per cent.

The average composition of five shales from the type Fram Fm is 65 ± 6 per cent quartz; 3 ± 1 per cent feldspar; 15 ± 4 per cent kaolinite; 12 ± 3 per cent illite; 1 ± 1 per cent chlorite; and 1 ± 1 per cent calcite.

Paleocurrents — Cross-beds were measured throughout the type section of the Fram Fm. The paleocurrent pattern (Fig. 30) is unimodal and indicates sediment transport to the southwest. Visual estimates of cross-bed orientation on eastern Grinnell Peninsula suggest a similar sediment transport direction.

Age — Palynological control is limited to spot samples collected near the base, middle and top of the formation of both the type section and the east Grinnell Peninsula locality. The available data indicate an age of Early to Middle Frasnian (*Medius* and *Maclarenii* zones) (Chi, *pers. comm.*, 1975). Plant and fish collections made by the Fram expedition also indicate a Late Devonian age (Nathorst, 1904; Kiaer, 1915).

Environment of Deposition — The lithologies, sedimentary structures, cyclicity and fossil content of the Fram Fm suggest a meandering-stream environment of deposition. Sandstone units are interpreted to be point-bar and channel-fill deposits, whereas the shale-siltstone units are of overbank origin.

The Fram Fm is similar to the Catskill Fm of Pennsylvania (*pers. obs.*), which is also interpreted to be of meandering-stream origin (Allen and Friend, 1968). The main lithologic difference between the two is

---

PLATE 24

Fig. 1. Basal conglomeratic portion of sandstone unit in Fram Fm, Bird Fiord area. Note basal scour surface and plant fragments (upper right) in sandstone.

Fig. 2. Siltstone with root markings, Fram Fm, Bird Fiord area.

Fig. 3. Siltstone with mud cracks, Fram Fm, Bird Fiord area.

Fig. 4. Photomicrograph of fine-grained, quartzose lithic arenite from Fram Fm, Bird Fiord area.

Composition — Fine-grained sandstones in the Fram Fm are quartz arenites and quartzose lithic arenites (Plate 24, Fig. 4). The average composition of five fine-grained sandstones from the type section is 70 ± 3 per cent quartz; 1 ± 1 per cent feldspar; 5 ± 2 per cent rock fragments (mainly shale); 6 ± 2 per cent chert; 12 ± 3 per cent clay (mainly kaolinite, minor illite and chlorite); 3 ± 3 per cent calcite; 2 ± 2 per cent porosity. Biotite and muscovite are also present and are usually concentrated in dark laminations along with carbonaceous material. Heavy minerals are rare and consist mainly of tourmaline, zircon, rutile and garnet. Most grains are subround.

Part of the quartz content is authigenic cement. The clay is interpreted to be mainly of diagenetic origin and tends to occur in the coarser

Age — Palynological control is limited to spot samples collected near the base, middle and top of the formation of both the type section and the east Grinnell Peninsula locality. **The available data indicate an age of Early to Middle Frasnian** (*Medius* and *Maclarenii* zones) (Chi, *pers. comm.*, 1975). Plant and fish collections made by the Fram expedition also indicate a Late Devonian age (Nathorst, 1904; Kiaer, 1915).

transport direction.

Age — Palynological control is limited to spot samples collected near the base, middle and top of the formation of both the type section and the east Grinnell Peninsula locality. The available data indicate an age of Early to Middle Frasnian (*Medius* and *Maclarenii* zones) (Chi, *pers. comm.*, 1975). Plant and fish collections made by the Fram expedition also indicate a Late Devonian age (Nathorst, 1904; Kiaer, 1915).

Environment of Deposition — The lithologies, sedimentary structures, cyclicity and fossil content of the Fram Fm suggest a meandering-stream environment of deposition. Sandstone units are interpreted to be point-bar and channel-fill deposits, whereas the shale-siltstone units are of overbank origin.

The Fram Fm is similar to the Catskill Fm of Pennsylvania (*pers. obs.*), which is also interpreted to be of meandering-stream origin (Allen and Friend, 1968). The main lithologic difference between the two is

PLATE 24

Fig. 1. Basal conglomeratic portion of sandstone unit in Fram Fm, Bird Fiord area. Note basal scour surface and plant fragments (upper right) in sandstone.

Fig. 2. Siltstone with root markings, Fram Fm, Bird Fiord area.

Fig. 3. Siltstone with mud cracks, Fram Fm, Bird Fiord area.

Fig. 4. Photomicrograph of fine-grained, quartzose lithic arenite from Fram Fm, Bird Fiord area.

Composition — Fine-grained sandstones in the Fram Fm are quartz arenites and quartzose lithic arenites (Plate 24, Fig. 4). The average composition of five fine-grained sandstones from the type section is 70 ± 3 per cent quartz; 1 ± 1 per cent feldspar; 5 ± 2 per cent rock fragments (mainly shale); 6 ± 2 per cent chert; 12 ± 3 per cent clay (mainly kaolinite, minor illite and chlorite); 3 ± 3 per cent calcite; 2 ± 2 per cent porosity. Biotite and muscovite are also present and are usually concentrated in dark laminations along with carbonaceous material. Heavy minerals are rare and consist mainly of tourmaline, zircon, rutile and garnet. Most grains are subround.

Part of the quartz content is authigenic cement. The clay is interpreted to be mainly of diagenetic origin and tends to occur in the coarser

Age — Palynological control is limited to spot samples collected near the base, middle and top of the formation of both the type section and the east Grinnell Peninsula locality. **The available data indicate an age of Early to Middle Frasnian** (*Medius* and *Maclarenii* zones) (Chi, *pers. comm.*, 1975). Plant and fish collections made by the Fram expedition also indicate a Late Devonian age (Nathorst, 1904; Kiaer, 1915).

transport direction.

Age — Palynological control is limited to spot samples collected near the base, middle and top of the formation of both the type section and the east Grinnell Peninsula locality. The available data indicate an age of Early to Middle Frasnian (*Medius* and *Maclarenii* zones) (Chi, *pers. comm.*, 1975). Plant and fish collections made by the Fram expedition also indicate a Late Devonian age (Nathorst, 1904; Kiaer, 1915).

Environment of Deposition — The lithologies, sedimentary structures, cyclicity and fossil content of the Fram Fm suggest a meandering-stream environment of deposition. Sandstone units are interpreted to be point-bar and channel-fill deposits, whereas the shale-siltstone units are of overbank origin.

The Fram Fm is similar to the Catskill Fm of Pennsylvania (*pers. obs.*), which is also interpreted to be of meandering-stream origin (Allen and Friend, 1968). The main lithologic difference between the two is

The Fram Fm is similar to the Catskill Fm of Pennsylvania (*pers. obs.*), which is also interpreted to be of meandering-stream origin (Allen and Friend, 1968). The main lithologic difference between the two is

Fiord area.



Middle-Upper Devonian at surface

1999 camps

2000 camps

2002 camps

ELLESMERE ISLAND

ARCTIC OCEAN

MELVILLE IS.

100km

AGE	SOUTHERN ELLESMERE ISLAND
DEVONIAN LATE FAMENNIAN	
DEVONIAN LATE FRASNIAN	NORDSTRAND POINT FORMATION
DEVONIAN LATE FRASNIAN	HELL GATE FM.
DEVONIAN LATE FRASNIAN	FRAM FM.
DEVONIAN MIDDLE GIVETIAN	HECLA BAY FORMATION
DEVONIAN MIDDLE EIFELIAN	STRATHCONA FIORD FM.
DEVONIAN MIDDLE EIFELIAN	UPPER BIRD FIORD FORMATION
DEVONIAN MIDDLE EIFELIAN	LOWER BIRD FIORD FORMATION


Middle-Upper Devonian clastic wedge at surface or in subsurface
1999 camps
2000 camps
2002 camps
2004 camp
100km
ELLESMERE ISLAND
AXEL HEIBERG
ELLEF RINGNES
AMUND RINGNES
MELVILLE
BATHURST
CORNWALLIS
DEVON I.
JONES SOUND
VISCOUNT MELVILLE SOUND
LANCASTER SOUND

Middle-Upper Devonian clastic wedge at surface or in subsurface
1999 camps
2000 camps
2002 camps
2004 camp
100km
ELLESMERE ISLAND
AXEL HEIBERG
ELLEF RINGNES
MELVILLE
BATHURST
DEVON I.
LANCASTER SOUND


Highlands
Tidal
Flats
Sea
O
SC
V
W
S

Middle-Upper Devonian clastic wedge at surface or in subsurface
1999 camps
2000 camps
2002 camps
2004 camp
100km
ELLESMERE ISLAND
AXEL HEIBERG
ELLEF RINGNES
MELVILLE
BATHURST
DEVON I.
VISCOUNT MELVILLE SOUND
LANCASTER SOUND

Middle-Upper Devonian clastic wedge at surface or in subsurface
1999 camps
2000 camps
2002 camps
2004 camp
100km
ELLESMERE
ISLAND
AXEL HEIBERG
ELLEF RINGNES
MELVILLE
BATHURST
CORNWALLIS
DEVON I.
VISCOUNT MELVILLE SOUND

Site NV2K17, southern Ellesmere Island, 2006



modified from Radinsky 1987






a
b

Heather King








COM
PF


a
possible fore fin contact
probable fore fin contact
hind fin contact
b
c





MID to LATE DEVONIAN



Developmental genetics of non-model organisms

Expeditions targeted to key parts of the tree of life

Early Hox Phases

d13	d12	d11	d10	d9	d8



Early Hox Phases

d13 | d12 | d11 | d10 | d9 | d8





Late Hox Phase



Origin of Late Hox Phases?


mtr4
dr
lep
app

Polyodon pectoral fins - *in situ* hybridization of 5' HoxD genes

Hoxd11

Hoxd12

Hoxd13

Early

Late

Polyodon pectoral fins - *in situ* hybridization of 5' HoxD genes



Early

Late

Polyodon pectoral fins - *in situ* hybridization of 5' HoxD genes





Source: Deschamps, 2004


Late
Phase
Early
Phase


Late
Phase
Early
Phase


GCR
HoxD cluster
CsB
CsA
Lnp
CsC
Evx2
350 kbp


Late
Phase
Early
Phase
CsB
CsA
GCR
HoxD cluster
CsB
CsA
Lnp
CsC
Evx2
350 kbp



Modified from: Spitz & Duboule, 2003



Modified from: Spitz & Duboule, 2003



Modified from: Spitz & Duboule, 2003


Late
Phase
Early
Phase
CsB
CsA
GCR
HoxD cluster
CsB
CsA
Lnp
CsC
Evx2
350 kbp
2.6 kbp
CsB
Mouse
Chicken
Frog
Zebrafish
Skate


GCR
HoxD cluster →
CsB
CsA
← Lnp
CsC
← Evx2
350 kbp


GCR
HoxD cluster →
CsB
CsA
← Lnp
CsC
← Evx2
350 kbp


LacZ
CsB
β globin -
LacZ


GCR
HoxD cluster →
CsB
CsA
← Lnp
CsC
← Evx2
350 kbp
CsB
LacZ
β globin -
LacZ


GCR
HoxD cluster →
CsB
CsA
← Lnp
CsC
← Evx2
350 kbp
CsB
LacZ
β globin -
LacZ
CsB

Chicken CsB



Chicken CsB



Skate CsB






Chondrichthyans
Danio rerio
Sauripterus taylori
Polyodon spathulae
Eusthenopteron foordi
Panderichthys rhombolepis
2008 reconstruction
Tiktaalik roseae
Acanthostega gunnari
Tulerpeton curtum
Gallus gallus
Mus musculus


Day 4
Day 5


AER
Growth
Pattern
Gre
BMP
Fgf
Fmn
Shh
ZPA

How did vertebrate appendages arise?



How did vertebrate appendages arise?
How do fins differ from other outgrowths?



mares

Shark gill skeleton



Little skate (*Leucoraja erinacea*)

Shark gill skeleton

Little skate (*Leucoraja erinacea*)

Shark gill skeleton



Little skate (*Leucoraja erinacea*)

“AER”

chick limb

AER



“AER”

chick limb

skate fin



AER



AEF



“AER”

chick limb

AER



skate fin



AEF



skate arch

‘AER’



“ZPA”



chick limb



skate fin

“ZPA”



chick limb



skate fin



skate arches

“ZPA”



chick limb



skate fin



skate arches

Wildtype

RA treatment

chick limb



Chick limb

Wildtype



RA treatment



Wildtype

RA treatment

Chick limb





Skate arch





Mirror-image, supernumerary rays

Wildtype

RA treatment

Chick limb





Wildtype

RA

Skate arch









Mirror-image, supernumerary rays


Gill
Fin
pr
rad
mt
Limb
digits
ra
hu
ul


Gill
Fin
Limb
Gill
Fin
Limb
pr
rad
mt
digits
ra
hu
ul


Shh
Fgf8
Fin/Limb
Arch
Fin/Limb
Arch
Skate
Chick
II
III
IV
II
III
IV
Gill
Fin
Limb
Gill
Fin
Limb
pr
rad
mt
digits
ra
hu
ul

Developmental genetics of non-model organisms

Expeditions targeted to key parts of the tree of life

EXTANT

EXTINCT




A
B
C
D
E
F
G
H
I
J
K
L

There are: unknown unknowns (sensu RUMSFELD, 2002)




Arctic Ocean
GREENLAND
Baffin Bay
Devon Island
Princess Royal Island
Bruce Pt
Bere Bay
Wilmer Bay
Tucker Pt
Cape Disraeli
Cape Separation
Discovery Mountain
Arthur Fjord
Q
OE
OCB
DBi
CPBC
DHB
DF
DHG
DNP
DPI
BERE BAY SYNCLINE
PARKER FAULT ZONE
MOUNT WILMER FAULT
TUCKER ANTICLINE
ARTHUR FJORD SYNCLINE
GRINNELL RANGE THRUST
MOUNT DAUNTLESS FAULT
NOTE 3
HADDINGTON

- Igor Schneider
- Randy Dahn
- Marcus Davis
- Andrew Gillis
- Justin Lemberg

- Igor Schneider
- Randy Dahn
- Marcus Davis
- Andrew Gillis
- Justin Lemberg
- Ted Daeschler
- Farish A. Jenkins, Jr.
- Jason Downs

- Igor Schneider
- Randy Dahn
- Marcus Davis
- Andrew Gillis
- Justin Lemberg

- Ted Daeschler
- Farish A. Jenkins, Jr.
- Jason Downs

- Marcelo Nobriega, Ivy Aneas
- Melina Hale, Heather King

- Igor Schneider
- Randy Dahn
- Marcus Davis
- Andrew Gillis
- Justin Lemberg

- Marcelo Nobriega, Ivy Aneas
- Melina Hale, Heather King

- Ted Daeschler
- Farish A. Jenkins, Jr.
- Jason Downs

- Nunavut Territory
- Polar Shelf Project
- Grise Fiord Community

- Igor Schneider
- Randy Dahn
- Marcus Davis
- Andrew Gillis
- Justin Lemberg
- Marcelo Nobriega, Ivy Aneas
- Melina Hale, Heather King

- Ted Daeschler
- Farish A. Jenkins, Jr.
- Jason Downs
- Nunavut Territory
- Polar Shelf Project
- Grise Fiord Community

- NSF, NGS, NIH, The University of Chicago
- Academy of Natural Sciences, MCZ Harvard